

Neurobiology of Speech Production

Pascale Tremblay^{1,2}, Isabelle Deschamps^{1,2} and Vincent L. Gracco^{3,4,5}

¹Centre de Recherche de l'Institut Universitaire en Santé Mentale de Québec, Québec City, QC, Canada; ²Département de Réadaptation, Faculté de Médecine, Université Laval, Québec City, QC, Canada; ³Centre for Research on Brain, Language and Music, McGill University, Montreal, QC, Canada; ⁴School of Communication Sciences and Disorders, McGill University, Montreal, QC, Canada; ⁵Haskins Laboratories, New Haven, CT, USA

59.1 INTRODUCTION

s0010

p0010

Speech is one of the most distinguishing human traits. It represents a model neural system for studying a range of human characteristics from sensorimotor control to cognition. It uses a complex control system optimized for sequential output and is used for both self-expressive and interactive communication. The production of speech reflects a complex and dynamic process dependent on the interaction among multiple cortical and subcortical regions for the fine control of more than 100 muscles located in the oral cavity, neck, and abdomen (see [Figure 59.1](#) for an overview). In the following, we identify the set of processes involved in speech production as well as their neural substrate.

from studies using speech errors. Speech errors can provide valuable insights regarding the linguistic principles that are involved in the production of speech because these errors are generally consistent with language-specific phonological rules ([Goldrick & Daland, 2009](#)). Based on speech error analyses, some researchers have postulated that the units of speech planning are individual phonological features ([Mowrey & MacKay, 1990](#)), whereas others propose that these units are bigger [e.g., phonemes ([Roelofs, 1997, 1999](#); [Stemberger, 1982](#)), syllables ([Levelt, 1999](#)), or words]. Recently, Peeva and colleagues ([2010](#)) used functional magnetic resonance imaging (fMRI) and a repetition–suppression (RS) paradigm ([Grill-Spector, Henson, & Martin, 2006](#); [Grill-Spector & Malach, 2001](#)) to study speech representations. Capitalizing on the RS phenomenon in which the repetition of a stimulus leads to reduced neural activity ([Henson & Rugg, 2003](#)), the authors varied the repetition rate of phonemes, syllables, and pseudowords expecting that areas sensitive to the processing of a specific type of phonological unit would show specific RS effects. Sensitivity to phonemic information was found in the supplementary motor area (SMA), the left palladium (in the basal ganglia or BG), the left posterior superior temporal gyrus (STG), and the left superior posterior lateral cerebellum. Sensitivity to syllable level information was found in the ventral premotor cortex (vPMC), which was also sensitive to phonemic information. Finally, sensitivity to supra-syllabic information was found in the right superior posterior lateral cerebellum. These results suggest that multiple levels of representations, including phonemic and syllabic, are involved in the production of speech sounds.

s0015

59.2 NEUROBIOLOGY OF SPEECH MOTOR CONTROL

s0020

59.2.1 Speech Representations

p0015

From a linguistic perspective, a number of potential candidate constructs may be represented in the neural processes associated with the production of speech. One view in the psycholinguistic literature is that grammatical encoding, or the creation of lexical items within a syntactic frame, and phonological encoding, including the specification of prosodic structure, are the two fundamental processes that create the phonetic plan ([Garrett, 1993](#); [Levelt, 1992, 1993](#)). The phonetic plan interfaces seamlessly with speech motor processes that generate the sequence of sounds specified in the plan.

p0020

One approach to associate these broad psycholinguistic processes with their neural substrates comes




FIGURE 59.1 Simplified illustration of the motor speech system including main cortical components and their connectivity, subcortical loops (cortico-striatal-thalamic loops and cortico-cerebellar-thalamic loops), as well as peripheral components (cranial and spinal nerve innervations to speech-related structures in the face, oral cavity, neck, and abdomen).

s0025 59.2.2 Speech Motor Planning and Programming

p0025 During speech production, phonological encoding is the retrieval of the phonological code that consists of segmental (phonemes, syllables) and suprasegmental information (such as stress). This information is used to build a representation of the syllabified word form. The syllabified word form provides the framework for the planning of a motor act. Although models of speech production converge on the notion that the output of phonological encoding is a phonological word in which metrical, syllabic, and segmental properties are fully specified, models of speech production differ with regard to whether the retrieval of the phonological code is presyllabified. For some, the syllabification is computed online depending on the context (Levelt, Roelofs, & Meyer, 1999), whereas for others the phonological code is presyllabified (Dell, 1988). Regardless of the theoretical perspective, phonological encoding is associated with the process of speech motor preparation, which involves the activation and translation of phonological representations into multiple domain-general mechanisms, including response selection, response sequencing, and movement initiation. These important mechanisms, often referred to

as “supra-motor functions” or “motor cognition” (Freund, Jeannerod, Hallett, & Leiguarda, 2005), are not specific to speech production but instead are part of the planning of all voluntary actions. Speech production builds on common action control mechanisms consistent with the notion that the speech system is an overlaid functional system that “[...] gets what service it can out of organs and functions, nervous and muscular, that have come into being and are maintained for very different ends than its own” (Sapir, 1921).

59.2.2.1 Response Selection

s0030
p0030 Response selection in spoken language production is the process by which a set of lexical units forming a message is transformed into motor programs, that is, stored motor routines. Several neuroimaging studies have examined the process of selecting nonspeech motor responses (such as finger and hand movements) and revealed activation in the presupplementary motor area (pre-SMA) (Brodmann’s medial area 6) in which the increase in activation is commensurate with demands on response selection. For instance, activation in pre-SMA is enhanced when participants are free to choose a motor response from among several alternatives compared with when they are required to execute a specific,

stimulus-driven, motor response (Deiber, Ibanez, Sadato, & Hallett, 1996; Lau, Rogers, & Passingham, 2006; Weeks, Honda, Catalan, & Hallett, 2001). Consistent with the nonspeech literature, several fMRI studies have shown that manipulating response selection during single word production modulates distributed brain networks including the pre-SMA, but also the adjacent cingulate motor area (CMA) and the vPMC (Crosson et al., 2001; Nagel, Schumacher, Goebel, & D'Esposito, 2008; Tremblay & Gracco, 2006; Tremblay & Small, 2011). Importantly, the pre-SMA is involved in selecting single words (Alario, Chainay, Lehericy, & Cohen, 2006; Tremblay & Gracco, 2006) but also noncommunicative oral motor gestures (Braun, Guillemin, Hosey, & Varga, 2001; Tremblay & Gracco, 2010), revealing a domain-general selection mechanisms. Moreover, transcranial magnetic stimulation (TMS) to the pre-SMA leads to impaired voluntary selection of actions, including words and noncommunicative oral motor gestures (Tremblay & Gracco, 2009), supporting the notion of a domain-general selection process. Taken together, these results suggest that the pre-SMA plays a central role in selecting motor responses for speech production. The pre-SMA has a connectivity pattern that is ideal for linking higher-level cognitive (including linguistic) and motor processes, a *sine qua non* for the implementation of response selection, with important projections from the prefrontal cortex, particularly the dorsolateral prefrontal cortex or DLPFC (Lu, Preston, & Strick, 1994; Luppino, Matelli, Camarda, & Rizzolatti, 1993), and connections with several nonprimary motor areas, such as the SMA-proper and the PMC (Luppino & Rizzolatti, 2000), for controlling motor output. Recent fMRI evidence suggests a role for the caudate nucleus in response selection for speech production (Argyropoulos, Tremblay, & Small, 2013), consistent with evidence on the anatomical connectivity of the caudate, which connects with the prefrontal as well as the SMA/pre-SMA (Di Martino et al., 2008; Lehericy et al., 2004), suggesting that response selection is implemented through cortico-striatal connections between the pre-SMA and the caudate nucleus.

s0035 59.2.2.2 Response Sequencing

p0035 In his classic article on serial order, Lashley described the problem of organizing component parts of an action into movement sequences as the *action syntax problem* (Lashley, 1951). The manifestation of action syntax can be seen in a multitude of behaviors ranging from human thought (Marsden, 1984) to grooming behavior in rats (Aldridge, Berridge, & Rosen, 2004). For speech, action sequences can be organized around multiple components (phonemes, syllables, words, phrases, etc.); without appropriate timing, in terms of either initiating the action or sequencing the action units, communication would be difficult. fMRI studies have shown that motor

sequencing is implemented in a network of regions organized around nonprimary motor areas (SMA-proper, PM), the cerebellum, and the BG (Bengtsson, Ehrsson, Forssberg, & Ullen, 2005; Gerloff, Corwell, Chen, Hallett, & Cohen, 1997; Macar et al., 2002). Repetitive TMS of the SMA-proper results in sequential timing disruptions in a complex finger movement task (Gerloff et al., 1997); SMA-proper activation accompanies tasks requiring the processing of temporal patterns (Macar et al., 2002). Using fMRI, Bohland and Guenther (2006) showed a bilateral network including the SMA, the anterior insula, and the superior cerebellum that was more strongly recruited for the production of complex sequences of syllables (ka-ru-ti) compared with the production of simpler sequences, which consisted of repeating the same sound three times (ta-ta-ta), consistent with the nonspeech literature. Although there remains a number of issues regarding the implementation of selection and sequencing mechanisms for speech, the available empirical evidence, though limited, suggests that speech production relies on common action control mechanisms centered on the pre-SMA, SMA, and vPMC.

59.2.2.3 Motor Programming

s0040

p0040 Preparing speech production also involves fine-tuning of the planned motor routines, including adjustments of velocity, muscle tone, movement range, and direction. Motor programming is necessary because even though a closed set of syllables is available in each language and probably stored as set of motor routines, syllables and words are never produced identically, they are co-articulated and modulated as a function of the linguistic, environmental, emotional, and social contexts. Motor programming is usually believed to involve both online feedback-based and feedforward control systems. According to Van der Merwe (2009), regions involved in programming include the cerebellum, SMA-proper, M1, and the BG, but experimental evidence is lacking. The issue of feedback-based motor control is discussed in Section 59.3.

59.2.2.4 Movement Initiation

s0045

p0045 The initiation and termination of an action is fundamental to all voluntary motor behaviors. For speech production, starting and stopping speech movements are associated with a diverse range of communicative actions such as turn-taking, producing a list of words, and the insertion of pauses for emphasis. The SMA-proper has been previously identified as contributing to speech timing (Brendel et al., 2010; Gracco, 1997) and, more generally, has been associated with sequence timing as well as the perception of time. Recently, using functional connectivity analysis, and evaluating the temporal dynamics of the BOLD signal, two separately organized networks for speech

production have been proposed (Brendel et al., 2010; Riecker et al., 2005) with the SMA and the insula identified as network components contributing to motor timing. For example, TMS to the SMA results in sequential timing disruptions (Gerloff et al., 1997), varying rate of stimulus presentation during reading results in modulation of SMA-proper activity (Price, Moore, Humphreys, Frackowiak, & Friston, 1996), and SMA activity accompanies tasks requiring the processing of temporal patterns (Bengtsson et al., 2005; Macar et al., 2002). Similarly, damage to the insula results in speech initiation impairments (Shuren, 1993) and apraxia of speech (Dronkers, 1996), a disorder of temporal sequencing, although there is controversy surrounding the role of insula in motor sequencing (Hillis et al., 2004). One possibility is that the SMA-proper and insula may be working to coordinate and time sequential actions, possibly through priming and then triggering motor cortex output. Speech and oral movements are localized around the central sulcus of the insula in an area that does not have direct projections onto lower motor neurons but does connect to frontal regions, including the DLPFC as well as the SMA and the sensorimotor portions of the striatum (Augustine, 1996). The DLPFC on the right hemisphere is known to modulate lower level systems (Shallice, 2004) and activity in the right DLPFC may be contributing to speech timing and/or temporal processing of action sequences (Coull, Frackowiak, & Frith, 1998; Vallesi, Shallice, & Walsh, 2007). It appears that the insula and SMA-proper form an integrated network component, operating in concert with peripheral feedback systems, to time sequential speech motor output. When there is a need for explicit timing control, prefrontal cortex participation is recruited.

Another way to study movement initiation is to compare the manner in which movements are triggered, whether externally by sensory events or at will. Movements initiated by external stimuli produce reliable activity in SMA-proper (Lee, Chang, & Roh, 1999; Thickbroom et al., 2000; Wiese et al., 2004) as well as in the left dorsal PMC (Krams, Rushworth, Deiber, Frackowiak, & Passingham, 1998; Lepage et al., 1999; Weeks et al., 2001), suggesting that these areas are involved in initiating actions based on external sensory triggers. In humans, TMS to the left PMC results in a response delay and a disruption in the early stage of reaching and grasping (before movement execution), suggesting a role in the onset of movement (Schluter, Rushworth, Passingham, & Mills, 1998). Importantly, the contrast of self-initiated and externally triggered movements reveals activation in the pre-SMA (Deiber et al., 1996; Jenkins, Jahanshahi, Jueptner, Passingham, & Brooks, 2000; Tsujimoto, Ogawa, Tsukada, Kakiuchi, & Sasaki, 1998), suggesting a role for this region in the

generation of an internal trigger to move, which supports a role for the SMA/pre-SMA in the timing and initiation of actions. Another potentially important region for the timing of actions and the generation of a movement trigger is the BG. In patients with Parkinson's disease (PD), a disorder of BG, there is a clear decline in the ability to initiate movements at will without a concomitant reduction or slowing of externally triggered actions (Cunnington, Ianssek, & Bradshaw, 1999; Freeman, Cody, & Schady, 1993; Praamstra, Stegeman, Cools, Meyer, & Horstink, 1998). Other BG dysfunctions lead to difficulty starting, stopping, or sustaining movements including speech (Speedie, Wertman, Ta'ir, & Heilman, 1993), as well as abnormal rate, regularity, and temporal ordering of speech movements (Ludlow, Connor, & Bassich, 1987; Skodda, 2011; Skodda & Schlegel, 2008; Volkmann, Hefter, Lange, & Freund, 1992), demonstrating the importance of BG for the timing of speech actions.

59.3 SPEECH MOVEMENT EXECUTION

s0050

The final output for speech comes mainly from the ventral part of the primary motor cortex (vM1), which contains the neurons controlling the vocal tract (Penfield & Boldrey, 1937). It has been estimated that approximately 100 striated and visceral muscles, distributed across the abdomen, neck, larynx, pharynx, and oral cavity, are involved in the production of speech, reflecting the immense complexity of this functional system, which, in mature speakers, may produce as many as 14 phonemes per second (i.e., between six and nine syllables per second) (Kent, 2000).

p0055

The pyramidal system, which includes the corticospinal and corticobulbar tracts, is one of the most important efferent pathways for the control of voluntary muscle contractions. It connects neurons in the cortex (upper motor neurons, UMN) to alpha (lower) motor neurons (LMN) located in the brainstem and spinal cord. LMN innervate the muscle fibers located in the face, neck, and abdomen. M1 is the cortical area that contains the largest number of pyramidal fibers (Kuypers, 1973; Murray & Coulter, 1981; Ralston & Ralston, 1985), particularly the giant Betz cells located in cortical layer V. However, anatomical studies have shown that M1 is connected through long and short association fibers to multiple nonprimary motor areas, including the SMA (Dum & Strick, 1991, 1996; Muakkassa & Strick, 1979) the CMA located just beneath the SMA on the dorsal and ventral banks of the cingulate sulcus (Dum & Strick, 1991; Muakkassa & Strick, 1979) and the dorsal and ventral PMC (Barbas & Pandya, 1987; Dum & Strick, 1991; He, Dum, & Strick, 1993). Importantly, these nonprimary motor areas

p0060

contain a high density of corticospinal and corticobulbar neurons, directly projecting to the spinal cord through the pyramidal tract (for a review of the connectivity of nonprimary motor areas, see [Picard & Strick, 1996, 2001](#)), and thus each has the potential to influence the generation and control of movement independently of M1 ([Dum & Strick, 1991](#)). Electrical stimulation of the SMA ([Fried et al., 1991](#); [Morris, Dinner, Luders, Wyllie, & Kramer, 1988](#); [Penfield & Welch, 1951](#); [Talairach & Bancaud, 1966](#)) and CMA ([von Cramon & Jurgens, 1983](#)) induce vocalization and speech arrests in humans, suggesting a role in the control of phonation and articulation for these regions.

p0065 The corticospinal tracts innervate motor nuclei located in the spinal cord, whereas corticobulbar fibers innervate motor nuclei located in the brainstem. Because the motor nuclei involved in the control of respiration (mainly expiration), phonation, and articulation are located in the pons, down to the lumbar portion of the spinal cord, the production of speech depends on the integrity of both the corticospinal tract, for the innervations of the muscles of respiration in the abdomen, neck, and shoulder, and the corticobulbar tract, for the sensorimotor innervations of laryngeal and supralaryngeal muscles (for reviews, see [Jurgens, 2002, 2009](#)) through six pairs of cranial nerves (CN V: trigeminal; CN VII: facial; CN IX: glossopharyngeal; CN X: vagus; CN XI: accessory; CN XII: hypoglossal).

p0070 All cortical axons (originating from M1, SMA, CMA, and PMC) forming the corticobulbar and corticospinal tracts converge into the internal capsule, located between the thalamus and BG, with fibers originating from ventral areas located rostrally to those originating from more dorsal areas ([Beever & Horsley, 1890](#); [Dejerine, 1901](#)). Most pyramidal fibers cross from one side to the other before entering the spinal cord at the level of the medulla oblongata (i.e., the pyramidal decussation); corticobulbar fibers cross at the level of the brainstem, although there are substantial bilateral innervations of the CN motor nuclei. The exceptions include contralateral innervations of ventral cell groups of the motor nucleus of the facial nerve (CN VII), which supply muscles of the lower quadrants of the face (e.g., the orbicularis oris muscle), and the hypoglossal nucleus (CN XII), which supplies the intrinsic and extrinsic muscles of the tongue.

p0075 It has been suggested that vocalizations are controlled through two distinct cortical-subcortical pathways, one involving a circuit formed by the CMA, the periaqueductal gray matter (PAG), and the reticular formation for the control of innate vocal patterns (e.g., crying, laughing, and moaning), and another connecting M1 to the phonatory motoneurons through the reticular formation for the control of patterned speech and singing ([Hsieh, Petrosyan, Goncalves, Hickok, & Saberi, 2011](#); [Jurgens,](#)

[2002](#)). This second circuit involves the cortico-striatal motor loop. Thus, M1 is connected not only to multiple nonprimary motor areas in the frontal lobe but also to the BG and cerebellum through the thalamus, and also to the reticular formation in the brainstem, controlling multiple aspects of speech production including respiration, vocalization, and articulation.

59.4 FEEDBACK PROCESSING AND SENSORY-MOTOR INTEGRATION

s0055

Early in the developmental process, the functional connection between speech perception and speech production is established and the ability to modify this coupling reflects the neural plasticity that continues throughout the life span. The resultant sensorimotor learning is the substrate on which developmental stages of speech and language develop, and one in which sensory feedback plays a crucial role ([Mowrer, 1952](#)). Somatosensory information from the lips and jaw have real-time access to modulate the spatial ([Abbs & Gracco, 1983](#); [Abbs, Gracco, & Cole, 1984](#); [Gracco & Abbs, 1985](#); [Shaiman & Gracco, 2002](#)) and temporal aspects of speech sequences ([Gracco & Abbs, 1989](#); [Saltzman, Lofqvist, Kay, Kinsella-Shaw, & Rubin, 1998](#)). Similar kinds of results are obtained from unanticipated alteration of auditory feedback for pitch ([Burnett, Freedland, Larson, & Hain, 1998](#); [Jones & Munhall, 2000](#)) and formants ([Purcell & Munhall, 2006a, 2006b](#)). The overarching conclusion is that sensory and motor systems for speech are in a constant state of interaction and integration and, most importantly, the sensorimotor integration forms the basis for successful and efficient speech production ([Gracco, 1991](#)).

p0080

From a control perspective, speech production can be conceptualized as representing a hybrid control scheme consisting of feedforward and feedback-like neural processes ([Abbs et al., 1984](#); [Guenther, Ghosh, & Tourville, 2006](#); [Hickok, 2012](#); [Houde & Nagarajan, 2011](#); [Tourville, Reilly, & Guenther, 2008](#)). Feedforward control is used to compute, before movement onset, the necessary motor commands that will achieve generally a desired movement goal given the system's current state. That is, the feedforward controller assembles a basic motor plan prior to movement onset and sends the commands to the appropriate musculature for execution. In contrast, feedback processes are used to adjust and correct motor commands that are planned or executed by the feedforward controller. However, if such adjustments would depend solely on afferent input signals, there would be an unavoidable delay that may be too long for movements as fast as those involved in many skilled actions, including speech production. Feedback control processes can also be used to

p0085

predict the sensory consequences of movements by making use of a copy of the prepared motor commands (efference copy or corollary discharge) (Sperry, 1950; Von Holst & Mittelstaedt, 1973). As such, information from somatosensory and auditory systems contributes in multiple ways. First, as part of the feedforward process, the sensory systems provide information about the initial conditions such that the motor commands for a desired outcome can be successfully achieved given the state of the vocal tract. Second, as part of the predictive process, they interact with the control signals to estimate the consequences of the planned action. Finally, as part of the feedback process, they modulate, in real time, adjustments to the motor commands based on re-afferent input during movement execution as well as signaling the achievement of the desired action.

p0090 The neural substrate associated with the sensorimotor aspects of speech production involves a mostly bilateral network of brain regions, including vM1 and sensory areas (somatosensory cortex, STG), nonprimary motor areas (vPMC, SMA-proper, CMA, and the insula), and subcortical regions associated with sensorimotor control (putamen, cerebellum, thalamus) (Ackermann & Riecker, 2004; Argyropoulos et al., 2013; Grabski, Tremblay, Gracco, Girin, & Sato, 2013; Riecker et al., 2004, 2005; Riecker, Wildgruber, Dogil, Grodd, & Ackermann, 2002; Tremblay, Deschamps, & Gracco, 2013; Tremblay & Gracco, 2006, 2009; Wise, Greene, Büchel, & Scott, 1999), areas that are known to receive afferent input from auditory and somatosensory areas. For example, in the macaque, the vPMC receives projection from sensory areas, including associative somatosensory area SII (Matelli, Camarda, Glickstein, & Rizzolatti, 1986) and the posterior STG (Chavis & Pandya, 1976; Schmahmann et al., 2007), whereas the SMA-proper receives important projection from the superior parietal lobule (area PEci, in the cingulate sulcus), which contains a complete somatosensory map of the body (Pandya & Seltzer, 1982) as well as from areas SII and SI (Luppino et al., 1993; McGuire, Bates, & Goldman-Rakic, 1991a, 1991b). Projections to the putamen have been reported from regions within the supratemporal plane and the STG (Yeterian & Pandya, 1998). The rostral and medial parts of STG project to rostroventral and caudoventral portions of the putamen, whereas the caudal portion of STG projects to caudal putamen. Recently, using resting state functional connectivity in humans, the dorsal portion of the putamen has been shown to connect with regions of the temporal cortex (Di Martino et al., 2008). As such, reafference may be an important source of information to assist in both the spatial and timing adjustments for the dynamic modulation of speech motor output as well as signaling successful achievement of speech motor goals (Gracco & Abbs, 1989).

For the cerebellum, bilateral posterior lobe activation p0095 in the vicinity of hemisphere lobule VI (Schmahmann et al., 1999) has been consistently reported during speech production (Ackermann, Mathiak, & Riecker, 2007; Riecker et al., 2002; Wise et al., 1999), most likely reflecting cortico-ponto-cerebellar projections from and to M1 (Kelly & Strick, 2003), possibly as part of an efference copy signal. A second area of activation on the inferior portion of the cerebellar hemisphere lobule VIIIA has been associated with auditory (Tourville et al., 2008) and somatosensory (Golfinopoulos et al., 2011) perturbations, as well as with rhythmic orofacial movements (Corfield et al., 1999) and sequencing non-meaningful syllables (Bohland & Guenther, 2006; Riecker, Kassubek, Groschel, Grodd, & Ackermann, 2006). The posterior lobe of the cerebellum receives sensory input from the trigeminal nerve (which provides sensorimotor innervations of the muscles of mastication) as well as the auditory system (Huerta, Frankfurter, & Harting, 1983; Ikeda & Matsushita, 1992; Pastor et al., 2002), and this area of the cerebellum may be implicated in multisensory rather than motor processing (Stoodley & Schmahmann, 2009; Thickbroom, Byrnes, & Mastaglia, 2003). Hence, sensory information from the dynamics of speech articulation has access to multiple brain regions through cortico-cortico, cortico-striatal, and cortico-cerebellar control loops.

59.5 CONCLUSION

s0060

In this chapter, we have shown that the neural system that controls speech production is immensely complex at all levels of the nervous system, involving multiple sensorimotor regions for motor planning and execution including M1, SMA, pre-SMA, CMA, the insula, and the supratemporal and inferior parietal cortices. Loops of internal control involving the BG, thalamus, and cerebellum are also involved in several aspects of speech movement preparation, including sequencing and temporal ordering. These regions work in concert to assemble complex, temporally ordered, and co-articulated sequences of speech movements; motor commands are sent through corticospinal and corticobulbar tracts involving seven cranial nerves, multiple spinal nerves, and more than 100 striatal and visceral nerves. Despite this remarkable complexity, the chain of events that leads to the production of speech occurs within several hundreds of milliseconds.

p0100

Acknowledgments

s0065

This project was funded by grants from the National Science and Engineering Research Council of Canada to PT and VLG, and from the National Institutes of Health Research (NIDCD—R01DC012502) to VLG. p0105

References

- Abbs, J. H., & Gracco, V. L. (1983). Sensorimotor actions in the control of multimovement speech gestures. *Trends in Neurosciences*, 6(9), 391–395.
- Abbs, J. H., Gracco, V. L., & Cole, K. J. (1984). Control of multimovement coordination: Sensorimotor mechanisms in speech motor programming. *Journal of Motor Behavior*, 16(2), 195–231.
- Ackermann, H., Mathiak, K., & Riecker, A. (2007). The contribution of the cerebellum to speech production and speech perception: Clinical and functional imaging data. *Cerebellum*, 6(3), 202–213. Available from: <http://dx.doi.org/10.1080/14734220701266742>.
- Ackermann, H., & Riecker, A. (2004). The contribution of the insula to motor aspects of speech production: A review and a hypothesis. *Brain and Language*, 89(2), 320–328. Available from: [http://dx.doi.org/10.1016/S0093-934X\(03\)00347-X](http://dx.doi.org/10.1016/S0093-934X(03)00347-X).
- Alario, F. X., Chainay, H., Lehericy, S., & Cohen, L. (2006). The role of the supplementary motor area (SMA) in word production. *Brain Research*, 1076(1), 129–143.
- Aldridge, J. W., Berridge, K. C., & Rosen, A. R. (2004). Basal ganglia neural mechanisms of natural movement sequences. *Canadian Journal of Physiology and Pharmacology*, 82(8–9), 732–739.
- Argyropoulos, G. P., Tremblay, P., & Small, S. L. (2013). The neostriatum and response selection in overt sentence production: An fMRI study. *NeuroImage*, 82C, 53–60. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2013.05.064>.
- Augustine, J. R. (1996). Circuitry and functional aspects of the insular lobe in primates including humans. *Brain Research Brain Research Reviews*, 22(3), 229–244.
- Barbas, H., & Pandya, D. N. (1987). Architecture and frontal cortical connections of the premotor cortex (area 6) in the rhesus monkey. *Journal of Comparative Neurology*, 256(2), 211–228.
- Beevor, C. E., & Horsley, V. (1890). An experimental investigation into the arrangement of the excitable fibres of the internal capsule of the Bonnet Monkey (*Macacus sinicus*). *Philosophical Transactions of the Royal Society of London B*, 181, 49–88.
- Bengtsson, S. L., Ehrsson, H. H., Forssberg, H., & Ullen, F. (2005). Effector-independent voluntary timing: Behavioural and neuroimaging evidence. *The European Journal of Neuroscience*, 22(12), 3255–3265. Available from: <http://dx.doi.org/10.1111/j.1460-9568.2005.04517.x>.
- Bohland, J. W., & Guenther, F. H. (2006). An fMRI investigation of syllable sequence production. *NeuroImage*, 32(2), 821–841. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2006.04.173>.
- Braun, A. R., Guillemin, A., Hosey, L., & Varga, M. (2001). The neural organization of discourse: An H2 15O-PET study of narrative production in English and American sign language. *Brain*, 124(10), 2028–2044.
- Brendel, B., Hertrich, I., Erb, M., Lindner, A., Riecker, A., Grodd, W., et al. (2010). The contribution of mesiofrontal cortex to the preparation and execution of repetitive syllable productions: An fMRI study. *NeuroImage*, 50(3), 1219–1230. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2010.01.039>.
- Burnett, T. A., Freedland, M. B., Larson, C. R., & Hain, T. C. (1998). Voice F0 responses to manipulations in pitch feedback. *The Journal of the Acoustical Society of America*, 103(6), 3153–3161.
- Chavis, D. A., & Pandya, D. N. (1976). Further observations on cortico-frontal connections in the rhesus monkey. *Brain Research*, 117(3), 369–386. Available from: [http://dx.doi.org/0006-8993\(76\)90089-5](http://dx.doi.org/0006-8993(76)90089-5) [pii].
- Corfield, D. R., Murphy, K., Josephs, O., Fink, G. R., Frackowiak, R. S., Guz, A., et al. (1999). Cortical and subcortical control of tongue movement in humans: A functional neuroimaging study using fMRI. *Journal of Applied Physiology*, 86(5), 1468–1477.
- Coull, J. T., Frackowiak, R. S., & Frith, C. D. (1998). Monitoring for target objects: Activation of right frontal and parietal cortices with increasing time on task. *Neuropsychologia*, 36(12), 1325–1334.
- Crosson, B., Sadek, J. R., Maron, L., Gökçay, D., Mohr, C., Auerbach, E. J., et al. (2001). Relative shift in activity from medial to lateral frontal cortex during internally versus externally guided word generation. *Journal of Cognitive Neuroscience*, 13(2), 272–283.
- Cunnington, R., Iansek, R., & Bradshaw, J. L. (1999). Movement-related potentials in Parkinson's disease: External cues and attentional strategies. *Movement Disorders*, 14(1), 63–68.
- Deiber, M. P., Ibanez, V., Sadato, N., & Hallett, M. (1996). Cerebral structures participating in motor preparation in humans: A positron emission tomography study. *Journal of Neurophysiology*, 75(1), 233–247.
- Dejerine, J. J. (1901). *Anatomie des centres nerveux* (Vol. 2). Paris: Rueff et Cie.
- Dell, G. S. (1988). The retrieval of phonological forms in production: Tests of predictions from a connectionist model. *Journal of Memory and Language*, 27(2), 124–142.
- Di Martino, A., Scheres, A., Margulies, D. S., Kelly, A. M., Uddin, L. Q., Shehzad, Z., et al. (2008). Functional connectivity of human striatum: A resting state fMRI study. *Cerebral Cortex*, 18(12), 2735–2747. Available from: <http://dx.doi.org/10.1093/cercor/bhn041>.
- Dronkers, N. F. (1996). A new brain region for coordinating speech articulation. *Nature*, 384(6605), 159–161. Available from: <http://dx.doi.org/10.1038/384159a0>.
- Dum, R. P., & Strick, P. L. (1991). The origin of corticospinal projections from the premotor areas in the frontal lobe. *Journal of Neuroscience*, 11(3), 667–689.
- Dum, R. P., & Strick, P. L. (1996). Spinal cord terminations of the medial wall motor areas in macaque monkeys. *The Journal of Neuroscience*, 16(20), 6513–6525.
- Freeman, J. S., Cody, F. W., & Schady, W. (1993). The influence of external timing cues upon the rhythm of voluntary movements in Parkinson's disease. *Journal of Neurology, Neurosurgery and Psychiatry*, 56(10), 1078–1084.
- Freund, H. J., Jeannerod, M., Hallett, M., & Leiguarda, R. (2005). *Higher-order motor disorders: From neuroanatomy and neurobiology to clinical neurology*. Oxford: Oxford University Press.
- Fried, I., Katz, A., McCarthy, G., Sass, K. J., Williamson, P., Spencer, S. S., et al. (1991). Functional organization of human supplementary motor cortex studied by electrical stimulation. *The Journal of Neuroscience*, 11(11), 3656–3666.
- Garrett, M. F. (1993). Errors and their relevance for models of language production. In G. Blanken, J. Dittman, H. Grim, J. Marshall, & C. Wallech (Eds.), *Linguistic disorders and pathologies* (pp. 69–96). Berlin: de Gruyter.
- Gerloff, C., Corwell, B., Chen, R., Hallett, M., & Cohen, L. G. (1997). Stimulation over the human supplementary motor area interferes with the organization of future elements in complex motor sequences. *Brain*, 120(Pt 9), 1587–1602.
- Goldrick, M., & Daland, R. (2009). Linking speech errors and phonological grammars: Insights from Harmonic Grammar networks. *Phonology*, 26(01), 147–185. Available from: <http://dx.doi.org/10.1017/S0952675709001742>.
- Golfinopoulos, E., Tourville, J. A., Bohland, J. W., Ghosh, S. S., Nieto-Castanon, A., & Guenther, F. H. (2011). fMRI investigation of unexpected somatosensory feedback perturbation during speech. *NeuroImage*, 55(3), 1324–1338. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2010.12.065>.
- Grabski, K., Tremblay, P., Gracco, V. L., Girin, L., & Sato, M. (2013). A mediating role of the auditory dorsal pathway in selective adaptation to speech: A state-dependent transcranial magnetic

- stimulation study. *Brain Research*, 1515, 55–65. Available from: <http://dx.doi.org/10.1016/j.brainres.2013.03.024>.
- Gracco, V. L. (1991). Sensorimotor mechanisms in speech motor control. In W. H. C. W. S. e. H. Peters (Ed.), *Speech motor control and stuttering* (pp. 53–78). Amsterdam: Elsevier.
- Gracco, V. L. (1997). A neuromotor perspective on speech production. In W. Hulstijn, H. F. M. Peters, & P. H. H. M. Van Lieshout (Eds.), *Speech production: Motor control, brain research and fluency disorders* (pp. 37–56). Amsterdam: Elsevier.
- Gracco, V. L., & Abbs, J. H. (1985). Dynamic control of the perioral system during speech: Kinematic analyses of autogenic and non-autogenic sensorimotor processes. *Journal of Neurophysiology*, 54(2), 418–432.
- Gracco, V. L., & Abbs, J. H. (1989). Sensorimotor characteristics of speech motor sequences. *Experimental Brain Research*, 75(3), 586–598.
- Grill-Spector, K., Henson, R., & Martin, A. (2006). Repetition and the brain: Neural models of stimulus-specific effects. *Trends in Cognitive Sciences*, 10(1), 14–23. Available from: <http://dx.doi.org/10.1016/j.tics.2005.11.006>.
- Grill-Spector, K., & Malach, R. (2001). fMR-adaptation: A tool for studying the functional properties of human cortical neurons. *Acta Psychologica*, 107(1–3), 293–321.
- Guenther, F. H., Ghosh, S. S., & Tourville, J. A. (2006). Neural modeling and imaging of the cortical interactions underlying syllable production. *Brain and Language*, 96(3), 280–301. Available from: <http://dx.doi.org/10.1016/j.bandl.2005.06.001>.
- He, S. Q., Dum, R. P., & Strick, P. L. (1993). Topographic organization of corticospinal projections from the frontal lobe: Motor areas on the lateral surface of the hemisphere. *The Journal of Neuroscience*, 13(3), 952–980.
- Henson, R. N., & Rugg, M. D. (2003). Neural response suppression, haemodynamic repetition effects, and behavioural priming. *Neuropsychologia*, 41(3), 263–270.
- Hickok, G. (2012). The cortical organization of speech processing: Feedback control and predictive coding the context of a dual-stream model. *Journal of Communication Disorders*. Available from: <http://dx.doi.org/10.1016/j.jcomdis.2012.06.004>.
- Hillis, A. E., Work, M., Barker, P. B., Jacobs, M. A., Breese, E. L., & Maurer, K. (2004). Re-examining the brain regions crucial for orchestrating speech articulation. *Brain*, 127(Pt 7), 1479–1487. Available from: [http://dx.doi.org/10.1093/brain/awh172awh172\[pil\]](http://dx.doi.org/10.1093/brain/awh172awh172[pil]).
- Houde, J. F., & Nagarajan, S. S. (2011). Speech production as state feedback control. *Frontiers in Human Neuroscience*, 5, 82. Available from: <http://dx.doi.org/10.3389/fnhum.2011.00082>.
- Hsieh, I. H., Petrosyan, A., Goncalves, O. F., Hickok, G., & Saberi, K. (2011). Observer weighting of interaural cues in positive and negative envelope slopes of amplitude-modulated waveforms. *Hearing Research*, 277(1–2), 143–151. Available from: <http://dx.doi.org/10.1016/j.heares.2011.01.008>.
- Huerta, M. F., Frankfurter, A., & Harting, J. K. (1983). Studies of the principal sensory and spinal trigeminal nuclei of the rat: Projections to the superior colliculus, inferior olive, and cerebellum. *The Journal of Comparative Neurology*, 220(2), 147–167. Available from: <http://dx.doi.org/10.1002/cne.902200204>.
- Ikeda, M., & Matsushita, M. (1992). Trigemino-cerebellar projections to the posterior lobe in the cat, as studied by anterograde transport of wheat germ agglutinin-horseradish peroxidase. *The Journal of Comparative Neurology*, 316, 221–237.
- Jenkins, I. H., Jahanshahi, M., Jueptner, M., Passingham, R. E., & Brooks, D. J. (2000). Self-Initiated versus externally triggered movements. II. The effect of movement predictability on regional cerebral blood flow. *Brain*, 123, 1216–1228.
- Jones, J. A., & Munhall, K. G. (2000). Perceptual calibration of F0 production: evidence from feedback perturbation. *The Journal of the Acoustical Society of America*, 108(3 Pt 1), 1246–1251.
- Jurgens, U. (2002). Neural pathways underlying vocal control. *Neuroscience and Biobehavioral Reviews*, 26(2), 235–258.
- Jurgens, U. (2009). The neural control of vocalization in mammals: A review. *Journal of Voice: Official Journal of the Voice Foundation*, 23(1), 1–10. Available from: <http://dx.doi.org/10.1016/j.jvoice.2007.07.005>.
- Kelly, R. M., & Strick, P. L. (2003). Cerebellar loops with motor cortex and prefrontal cortex of a nonhuman primate. *The Journal of Neuroscience*, 23(23), 8432–8444. Available from: <http://dx.doi.org/10.1523/JNEUROSCI.2323-03.2003> [pii].
- Kent, R. D. (2000). Research on speech motor control and its disorders: A review and prospective. *Journal of Communication Disorders*, 33(5), 391–427. quiz 428. Available from: [http://dx.doi.org/10.1016/S0021-9924\(00\)00023-X](http://dx.doi.org/10.1016/S0021-9924(00)00023-X) [pii].
- Krams, M., Rushworth, M. F., Deiber, M. P., Frackowiak, R. S., & Passingham, R. E. (1998). The preparation, execution and suppression of copied movements in the human brain. *Experimental Brain Research*, 120(3), 386–398.
- Kuypers, H. G. J. M. (1973). The anatomical organization of the descending pathways and their contribution to motor control especially in primates. In J. Desmedt (Ed.), *New developments in EMG and clinical neurophysiology* (pp. 38–68). Basel, Switzerland: Karger.
- Lashley, K. S. (1951). The problem of serial order in behavior. In L. A. Jeffress (Ed.), *Cerebral mechanisms in behavior* (pp. 112–131). New York, NY: Wiley.
- Lau, H., Rogers, R. D., & Passingham, R. E. (2006). Dissociating response selection and conflict in the medial frontal surface. *NeuroImage*, 29(2), 446–451.
- Lee, K. M., Chang, K. H., & Roh, J. K. (1999). Subregions within the supplementary motor area activated at different stages of movement preparation and execution. *NeuroImage*, 9, 117–123.
- Lehericy, S., Ducros, M., Krainik, A., Francois, C., Van de Moortele, P. F., Ugurbil, K., et al. (2004). 3-D diffusion tensor axonal tracking shows distinct SMA and pre-SMA projections to the human striatum. *Cerebral Cortex*, 14(12), 1302–1309. Available from: <http://dx.doi.org/10.1093/cercor/bhh091>.
- Lepage, M., Beaudoin, G., Boulet, C., O'Brien, I., Marcantoni, W., Bourgouin, P., et al. (1999). Frontal cortex and the programming of repetitive tapping movements in man: Lesion effects and functional neuroimaging. *Brain Research. Cognitive Brain Research*, 8(1), 17–25.
- Levelt, W. J. (1992). Accessing words in speech production: Stages, processes and representations. *Cognition*, 42(1–3), 1–22.
- Levelt, W. J. (1993). Timing in speech production with special reference to word form encoding. *Annals of the New York Academy of Sciences*, 682, 283–295.
- Levelt, W. J. (1999). Models of word production. *Trends in Cognitive Sciences*, 3(6), 223–232. Available from: [http://dx.doi.org/10.1016/S1364-6613\(99\)01319-4](http://dx.doi.org/10.1016/S1364-6613(99)01319-4) [pii].
- Levelt, W. J., Roelofs, A., & Meyer, A. S. (1999). A theory of lexical access in speech production. *The Behavioral and Brain Sciences*, 22(1), 1–38 (discussion 38–75).
- Lu, M. T., Preston, J. B., & Strick, P. L. (1994). Interconnections between the prefrontal cortex and the premotor areas in the frontal lobe. *Journal of Comparative Neurology*, 341(3), 375–392.
- Ludlow, C. L., Connor, N. P., & Bassich, C. J. (1987). Speech timing in Parkinson's and Huntington's disease. *Brain and Language*, 32(2), 195–214.
- Luppino, G., Matelli, M., Camarda, R., & Rizzolatti, G. (1993). Corticocortical connections of area F3 (SMA-proper) and area F6 (pre-SMA) in the macaque monkey. *Journal of Comparative Neurology*, 338(1), 114–140.
- Luppino, G., & Rizzolatti, G. (2000). The organization of the frontal motor cortex. *News and Views*, 15, 219–224.

- Macar, F., Lejeune, H., Bonnet, M., Ferrara, A., Pouthas, V., Vidal, F., et al. (2002). Activation of the supplementary motor area and of attentional networks during temporal processing. *Experimental Brain Research*, 142(4), 475–485. Available from: <http://dx.doi.org/10.1007/s00221-001-0953-0>.
- Marsden, C. D. (1984). Motor disorders in basal ganglia disease. *Human Neurobiology*, 2(4), 245–250.
- Matelli, M., Camarda, R., Glickstein, M., & Rizzolatti, G. (1986). Afferent and efferent projections of the inferior area 6 in the macaque monkey. *Journal of Comparative Neurology*, 251(3), 281–298.
- McGuire, P. K., Bates, J. F., & Goldman-Rakic, P. S. (1991a). Interhemispheric integration: I. Symmetry and convergence of the corticocortical connections of the left and the right principal sulcus (PS) and the left and the right supplementary motor area (SMA) in the rhesus monkey. *Cerebral Cortex*, 1(5), 390–407.
- McGuire, P. K., Bates, J. F., & Goldman-Rakic, P. S. (1991b). Interhemispheric integration: II. Symmetry and convergence of the corticostriatal projections of the left and the right principal sulcus (PS) and the left and the right supplementary motor area (SMA) of the rhesus monkey. *Cerebral Cortex*, 1(5), 408–417.
- Morris, H. H., III, Dinner, D. S., Luders, H., Wyllie, E., & Kramer, R. (1988). Supplementary motor seizures: Clinical and electroencephalographic findings. *Neurology*, 38(7), 1075–1082.
- Mowrer, O. H. (1952). Speech development in the young child. I. The autism theory of speech development and some clinical applications. *The Journal of Speech and Hearing Disorders*, 17(3), 263–268.
- Mowrey, R. A., & MacKay, I. R. (1990). Phonological primitives: Electromyographic speech error evidence. *The Journal of the Acoustical Society of America*, 88(3), 1299–1312.
- Muakkassa, K. F., & Strick, P. L. (1979). Frontal lobe inputs to primate motor cortex: Evidence for four somatotopically organized “premotor” areas. *Brain Research*, 177(1), 176–182.
- Murray, E. A., & Coulter, J. D. (1981). Organization of corticospinal neurons in the monkey. *The Journal of Comparative Neurology*, 195(2), 339–365. Available from: <http://dx.doi.org/10.1002/cne.901950212>.
- Nagel, I. E., Schumacher, E. H., Goebel, R., & D’Esposito, M. (2008). Functional MRI investigation of verbal selection mechanisms in lateral prefrontal cortex. *NeuroImage*, 43(4), 801–807. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2008.07.017>.
- Pandya, D. N., & Seltzer, B. (1982). Intrinsic connections and architectonics of posterior parietal cortex in the rhesus monkey. *The Journal of Comparative Neurology*, 204, 196–210.
- Pastor, M. A., Artieda, J., Arbizu, J., Marti-Clement, J. M., Penuelas, I., & Masdeu, J. C. (2002). Activation of human cerebral and cerebellar cortex by auditory stimulation at 40 Hz. *The Journal of Neuroscience*, 22(23), 10501–10506. Available from: <http://dx.doi.org/10.1523/JNEUROSCI.10501-02.2002>.
- Peeva, M. G., Guenther, F. H., Tourville, J. A., Nieto-Castanon, A., Anton, J. L., Nazarian, B., et al. (2010). Distinct representations of phonemes, syllables, and supra-syllabic sequences in the speech production network. *NeuroImage*, 50(2), 626–638. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2009.12.065>.
- Penfield, W., & Boldrey, E. (1937). Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain: A Journal of Neurology*, 60, 389–443.
- Penfield, W., & Welch, K. (1951). The supplementary motor area of the cerebral cortex; a clinical and experimental study. *A.M.A. Archives of Neurology and Psychiatry*, 66(3), 289–317.
- Picard, N., & Strick, P. L. (1996). Motor areas of the medial wall: A review of their location and functional activation. *Cerebral Cortex*, 6, 342–353.
- Picard, N., & Strick, P. L. (2001). Imaging the premotor areas. *Current Opinion in Neurobiology*, 11(6), 663–672.
- Praamstra, P., Stegeman, D. F., Cools, A. R., Meyer, A. S., & Horstink, M. W. (1998). Evidence for lateral premotor and parietal overactivity in Parkinson’s disease during sequential and bimanual movements. A PET study. *Brain*, 121, 769–772.
- Price, C. J., Moore, C. J., Humphreys, G. W., Frackowiak, R. S., & Friston, K. J. (1996). The neural regions sustaining object recognition and naming. *Proceedings Biological Sciences/The Royal Society*, 263(1376), 1501–1507. Available from: <http://dx.doi.org/10.1098/rspb.1996.0219>.
- Purcell, D. W., & Munhall, K. G. (2006a). Adaptive control of vowel formant frequency: Evidence from real-time formant manipulation. *The Journal of the Acoustical Society of America*, 120(2), 966–977.
- Purcell, D. W., & Munhall, K. G. (2006b). Compensation following real-time manipulation of formants in isolated vowels. *The Journal of the Acoustical Society of America*, 119(4), 2288–2297.
- Ralston, D. D., & Ralston, H. J., III (1985). The terminations of corticospinal tract axons in the macaque monkey. *The Journal of Comparative Neurology*, 242(3), 325–337. Available from: <http://dx.doi.org/10.1002/cne.902420303>.
- Riecker, A., Gerloff, C., Wildgruber, D., Nagele, T., Grodd, W., Dichgans, J., et al. (2004). Transient crossed aphasia during focal right-hemisphere seizure. *Neurology*, 63(10), 1932. Available from: <http://dx.doi.org/10.1213/00006123-200410000-00011> [pii].
- Riecker, A., Kassubek, J., Groschel, K., Grodd, W., & Ackermann, H. (2006). The cerebral control of speech tempo: Opposite relationship between speaking rate and BOLD signal changes at striatal and cerebellar structures. *NeuroImage*, 29(1), 46–53. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2005.03.046>.
- Riecker, A., Mathiak, K., Wildgruber, D., Erb, M., Hertrich, I., Grodd, W., et al. (2005). fMRI reveals two distinct cerebral networks subserving speech motor control. *Neurology*, 64, 700–706.
- Riecker, A., Wildgruber, D., Dogil, G., Grodd, W., & Ackermann, H. (2002). Hemispheric lateralization effects of rhythm implementation during syllable repetitions: An fMRI study. *NeuroImage*, 16, 169–176.
- Roelofs, A. (1997). The WEAVER model of word-form encoding in speech production. *Cognition*, 64(3), 249–284.
- Roelofs, A. (1999). Phonological segments and features as planning units in speech production. *Language and Cognitive Processes*, 14(2), 173–200. Available from: <http://dx.doi.org/10.1080/016909699386338>.
- Saltzman, E., Lofqvist, A., Kay, B., Kinsella-Shaw, J., & Rubin, P. (1998). Dynamics of intergestural timing: A perturbation study of lip-larynx coordination. *Experimental Brain Research*, 123(4), 412–424.
- Sapir, E. (1921). *Language: An introduction to the study of speech*. New York, NY: Harcourt, Brace and Company.
- Schluter, N. D., Rushworth, M. F., Passingham, R. E., & Mills, K. R. (1998). Temporary interference in human lateral premotor cortex suggests dominance for the selection of movements. A study using transcranial magnetic stimulation. *Brain*, 121(5), 785–799.
- Schmahmann, J. D., Doyon, J., McDonald, D., Holmes, C., Lavoie, K., Hurwitz, A. S., et al. (1999). Three-dimensional MRI atlas of the human cerebellum in proportional stereotaxic space. *NeuroImage*, 10(3 Pt 1), 233–260. Available from: <http://dx.doi.org/10.1006/nimg.1999.0459>.
- Schmahmann, J. D., Pandya, D. N., Wang, R., Dai, G., D’Arceuil, H. E., de Crespigny, A. J., et al. (2007). Association fibre pathways of the brain: Parallel observations from diffusion spectrum imaging and autoradiography. *Brain*, 130(Pt 3), 630–653. Available from: <http://dx.doi.org/10.1093/brain/awl359>.
- Shaiman, S., & Gracco, V. L. (2002). Task-specific sensorimotor interactions in speech production. *Experimental Brain Research*, 146(4), 411–418. Available from: <http://dx.doi.org/10.1007/s00221-002-1195-5>.

- Shallice, T. (2004). The fractionation of supervisory control. In M. S. Gazzaniga (Ed.), *The cognitive neurosciences* (pp. 943–956). Cambridge, MA: MIT Press.
- Shuren, J. (1993). Insula and aphasia. *Journal of Neurology*, 240(4), 216–218.
- Skodda, S. (2011). Aspects of speech rate and regularity in Parkinson's disease. *Journal of the Neurological Sciences*, 310(1-2), 231–236. Available from: <http://dx.doi.org/10.1016/j.jns.2011.07.020>.
- Skodda, S., & Schlegel, U. (2008). Speech rate and rhythm in Parkinson's disease. *Movement Disorders: Official Journal of the Movement Disorder Society*, 23(7), 985–992. Available from: <http://dx.doi.org/10.1002/mds.21996>.
- Speedie, L. J., Wertman, E., Ta'ir, J., & Heilman, K. M. (1993). Disruption of automatic speech following a right basal ganglia lesion. *Neurology*, 43(9), 1768–1774.
- Sperry, R. W. (1950). Neural basis of the spontaneous optokinetic response produced by visual inversion. *Journal of Comparative Physiology and Psychology*, 43, 482–489.
- Stemberger, J. P. (1982). The nature of segments in the lexicon: Evidence from speech errors. *Lingua*, 56(3–4), 235–259. Available from: [http://dx.doi.org/10.1016/0024-3841\(82\)90012-2](http://dx.doi.org/10.1016/0024-3841(82)90012-2).
- Stoodley, C. J., & Schmahmann, J. D. (2009). The cerebellum and language: Evidence from patients with cerebellar degeneration. *Brain and Language*, 110(3), 149–153. Available from: <http://dx.doi.org/10.1016/j.bandl.2009.07.006>.
- Talairach, J., & Bancaud, J. (1966). The supplementary motor area in man. *International Journal of Neurology*, 5330–5347.
- Thickbroom, G. W., Byrnes, M. L., & Mastaglia, F. L. (2003). Dual representation of the hand in the cerebellum: Activation with voluntary and passive finger movement. *NeuroImage*, 18(3), 670–674.
- Thickbroom, G. W., Byrnes, M. L., Sacco, P., Ghosh, S., Morris, I. T., & Mastaglia, F. L. (2000). The role of the supplementary motor area in externally timed movement: The influence of predictability of movement timing. *Brain Research*, 874(2), 233–241.
- Tourville, J. A., Reilly, K. J., & Guenther, F. H. (2008). Neural mechanisms underlying auditory feedback control of speech. *NeuroImage*, 39(3), 1429–1443. Available from: <http://dx.doi.org/10.1016/j.neuroimage.2007.09.054>.
- Tremblay, P., Deschamps, I., & Gracco, V. L. (2013). Regional heterogeneity in the processing and the production of speech in the human planum temporale. *Cortex; A Journal Devoted to the Study of the Nervous System and Behavior*, 49(1), 143–157. Available from: <http://dx.doi.org/10.1016/j.cortex.2011.09.004>.
- Tremblay, P., & Gracco, V. L. (2006). Contribution of the frontal lobe to externally and internally specified verbal responses: fMRI evidence. *NeuroImage*, 33(3), 947–957.
- Tremblay, P., & Gracco, V. L. (2009). Contribution of the pre-SMA to the production of words and non-speech oral motor gestures, as revealed by repetitive transcranial magnetic stimulation (rTMS). *Brain Research*, 1268, 112–124. Available from: <http://dx.doi.org/10.1016/j.brainres.2009.02.076>.
- Tremblay, P., & Gracco, V. L. (2010). On the selection of words and oral motor responses: Evidence of a response-independent frontoparietal network. *Cortex; A Journal Devoted to the Study of the Nervous System and Behavior*, 46(1), 15–28. Available from: <http://dx.doi.org/10.1016/j.cortex.2009.03.003>.
- Tremblay, P., & Small, S. L. (2011). Motor response selection in overt sentence production: A functional MRI study. *Frontiers in Psychology*, 2, 253. Available from: <http://dx.doi.org/10.3389/fpsyg.2011.00253>.
- Tsujimoto, T., Ogawa, M., Tsukada, H., Kakiuchi, T., & Sasaki, K. (1998). Activation of the ventral and mesial frontal cortex of the monkey by self-initiated movement tasks as revealed by positron emission tomography. *Neuroscience Letters*, 258(2), 117–120.
- Vallesi, A., Shallice, T., & Walsh, V. (2007). Role of the prefrontal cortex in the foreperiod effect: TMS evidence for dual mechanisms in temporal preparation. *Cerebral Cortex*, 17(2), 466–474. Available from: <http://dx.doi.org/10.1093/cercor/bhj163>.
- Van der Merwe, A. (2009). A theoretical framework for the characterization of pathological speech sensorimotor control. In M. R. McNeil (Ed.), *Clinical management of sensorimotor speech disorders* (pp. 3–18). New York, NY: Thieme.
- Volkman, J., Hefter, H., Lange, H. W., & Freund, H. J. (1992). Impairment of temporal organization of speech in basal ganglia diseases. *Brain and Language*, 43(3), 386–399. Available from: [http://dx.doi.org/10.1006/0093-934X\(92\)90108-Q](http://dx.doi.org/10.1006/0093-934X(92)90108-Q) [pii].
- von Cramon, D., & Jurgens, U. (1983). The anterior cingulate cortex and the phonatory control in monkey and man. *Neuroscience and Biobehavioral Reviews*, 7(3), 423–425.
- Von Holst, E., & Mittelstaedt, H. (1973). The reafference principle. In R. Martin (Ed.), *The behavioral physiology of animals and man. The collected papers of Erich von Holst* (Vol. 1, pp. 139–173). Coral Gables, FL: University of Miami Press.
- Weeks, R. A., Honda, M., Catalan, M. J., & Hallett, M. (2001). Comparison of auditory, somatosensory, and visually instructed and internally generated finger movements: A PET study. *NeuroImage*, 14(1), 219–230.
- Wiese, H., Stude, P., Nebel, K., de Greiff, A., Forsting, M., Diener, H. C., et al. (2004). Movement preparation in self-initiated versus externally triggered movements: An event-related fMRI-study. *Neuroscience Letters*, 371(2–3), 220–225.
- Wise, R. J., Greene, J., Büchel, C., & Scott, S. K. (1999). Brain regions involved in articulation. *Lancet*, 353, 1057–1061.
- Yeterian, E. H., & Pandya, D. N. (1998). Corticostriatal connections of the superior temporal region in rhesus monkeys. *The Journal of Comparative Neurology*, 399(3), 384–402.

NON-PRINT ITEM

Abstract

Speaking is a dynamic process in which neural representations associated with language and the constantly changing vocal tract configurations used to produce speech are seamlessly integrated to communicate. This integrative process requires a stable repertoire of motor routines and associated flexible sensorimotor and cognitive processes for speech preparation, execution, and monitoring. The development of neurobiologically plausible models of speech production requires an understanding of the role and contribution of both central and peripheral factors in the acquisition, maintenance, and reorganization of the brain-behavior patterns underlying speech production. Recent advances in brain imaging and neuromodulation methods, signal processing, and data analysis techniques have provided new insights into the organization of the brain networks and brain-behavior relations, leading to more comprehensive and more realistic models of speech production. In this chapter we identify the set of processes that are involved in producing speech and their neural substrate.

Keywords: Speech production; articulation; corticobulbar tract; sensorimotor integration; efference copy; speech planning; speech programming; cortico-striatal loops; nonprimary motor areas; cerebellum; basal ganglia; insula; supplementary motor area; premotor cortex; linguistic representations